

**DIPLOMA IN MEDICAL RECORD SCIENCE
SECOND YEAR
PAPER I – INTERNATIONAL CLASSIFICATION OF DISEASES (ICD-10) &
SURGICAL PROCEDURE (ICD-9CM)**

Q.P. Code: 841311

Time : Three Hours

Maximum : 100 Marks

Answer All questions.

I. Elaborate on:

(3 x 10 = 30)

1. a) What is the abbreviation for ICD-O?
b) What is the code number for unknown primary site?
c) Write two important situations the parenthesis is used in volume 1.
d) What is maternal death?
e) Write two conventions used in the Tabular list.

2. State whether the following statements are true or false:
a) Subdivision code '0' (zero) is used to indicate closed fracture.
b) Chapter IX deals with disease of the ear and mastoid process.
c) In dual coding system, the symbols "dagger and asterisk" are used.
d) Benign neoplasm code numbers are between (C00-C97).
e) Alphabet 'O' is used for pregnancy, child birth and the puerperium.

3. Fill in the blanks with suitable answers
(1) Neonatal period ends _____ completed days after birth.
(2) Post term is _____ completed weeks or more.
(3) _____ is the primary code for the underlying disease in the Dual coding system
(4) _____ is the abbreviation of NEC.
(5) Chapter XI refers the diseases of the _____ system.

II. Give the ICD 10 codes for the following:

(10 x 5 = 50)

1. a) Transitional cell carcinoma of bladder (write Morphology code also).
b) Congenital Stenosis of Urethra.

2. 25% with 2nd degree deep multiple site burn, exposure to accidental fire.

3. a) Postpartum Endometritis.
b) Adenotonsil Hypertrophy.

4. a) Medical Abortion Sepsis.
b) Infective Endocarditis.

5. a) Accessory breast
b) Achalasia Pylorus.

6. a) Sensorineural Bilateral hearing loss.
b) Dental caries.

7. a) Acromioclavicular joint dislocation.
b) Status Nephrectomy.
8. a) Acute Lymphoblastic Leukemia.
b) Systemic Lupus Erythematosus.
9. a) Non toxic multinodular Goitre.
b) Sequelae Fracture of femur.
10. a) Scrub typhus.
b) Motor neuron disease.

III. Give the ICD 9CM procedure codes for the following:

(10 x 2 = 20)

1. Partial Hypophysectomy trans-sphenoidal approach.
2. Wertheim operation.
3. Radical Mastoidectomy.
4. Take down Ileostomy.
5. Tap thorax.
6. Tenosynovectomy hand.
7. Epidural injection.
8. Excision Lesion skull.
9. Osteotomy femur wedge.
10. Axillary brachial vascular bypass graft.
